

IMPLICIT DIFFERENTIATION (Sec. 2.6)

Implicit differentiation is an application of chain rule. We need first to understand what means for a function to be implicitly defined.

Let us start from the definition of "explicit" and "implicit" that we can find in a dictionary.

EXPLICIT: stated clearly and in detail, leaving no room for confusion or doubt.

IMPLICIT: suggested but not communicated directly.

These two definitions give a very good idea of what means for a function to be explicitly or implicitly defined.

So far we have only met with functions explicitly defined. Indeed, when we write:

left side depends only on the dependent variable $y = f(x)$ right side depends only on the independent variable

we are defining **explicitly** our function. This means that the dependent variable is expressed explicitly in terms of the independent variable.

In this case we can easily compute the derivative $y' = f'(x)$ (in Lagrange notation) or $\frac{dy}{dx} = \frac{df}{dx}$ (in Leibniz notation) by applying the differentiation rules.

ex: $y = 3x^2 + \cos(x)$ is explicitly defined

$$\left. \begin{array}{l} \frac{dy}{dx} \\ y' \end{array} \right\} = 6x - \sin(x)$$

Now it is also possible to define a function implicitly as a function of x , through an equation that relates the independent and dependent variable.

Let us understand this on an example. Let us consider the equation:

$$x^2 + y^2 = 1$$

We will study this equation from two different points of view: algebraically and geometrically.

Algebraically

If we choose x as the independent variable, we say that y is implicitly defined as a function of x by the equation:

$$x^2 + y^2 = 1.$$

But attention: when y is implicitly defined, y is not a function in general.

Indeed if we fix an input for x there might exist several values of y that satisfy the equation, in other words several outputs.

ex: if $x=0 \Rightarrow 0^2 + y^2 = 1 \Rightarrow y^2 = 1 \Rightarrow y=1$ or $y=-1$

if $x = \frac{1}{2} \Rightarrow \left(\frac{1}{2}\right)^2 + y^2 = 1 \Rightarrow y^2 = \frac{3}{4} \Rightarrow y = \frac{\sqrt{3}}{2}$ or $y = -\frac{\sqrt{3}}{2}$

There are indeed two functions that are implicitly represented by the equation $x^2 + y^2 = 1$:

$$f_1(x) = \sqrt{1-x^2} \quad \text{and} \quad f_2(x) = -\sqrt{1-x^2}$$

Note: we say that $f(x)$ is a function implicitly defined by an equation $F(x,y)=0$ if $F(x, f(x))=0$

for all values x in the domain of f .

Indeed for f_1 and f_2 the equalities $x^2 + (f_1(x))^2 = 1$ and $x^2 + (f_2(x))^2 = 1$ are true for all values x in $[-1,1]$, which is the domain for both f_1 and f_2 .

The functions f_1 and f_2 can be found by solving the equation $x^2 + y^2 = 1$ for y .

$$x^2 + y^2 = 1 \Rightarrow y^2 = 1 - x^2 \Rightarrow y = \pm \sqrt{1 - x^2}.$$

Remark: Note that each explicit function can be written implicitly:

ex: $y = 3x^2 + \cos(x) \rightarrow \underbrace{y - 3x^2 - \cos(x)}_{F(x,y)} = 0$

Geometrically

An equation $F(x,y) = 0$ in the variables x and y corresponds in the real plane to a curve which is not in general the graph of a function.

The points of the curve are all the points of the plane whose coordinates (x,y) satisfy the equation $F(x,y) = 0$.


There exist also equations that correspond to curves without points

ex: $x^2 + y^2 = -1$ ← indeed the sum of two squares is always a non negative number


If we go back to our equation $x^2 + y^2 = 1$ we have that it corresponds to the circle of center $(0,0)$ and radius 1.

Indeed if $P(x,y)$ is a point in the plane, the quantity $x^2 + y^2$ represents its distance from the origin (Pythagorean theorem).

It is easy to see that the circle does not pass the vertical line test, so it is not the graph of a function.



However we can see the circle as the union of the graphs of the previous two functions f_1 and f_2



—————

Now that we have understood the difference between a function explicitly / implicitly defined, let us consider the following problem:

Problem: Find the tangent line to the circle $x^2 + y^2 = 1$ at the point $(\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2})$.

Solution: So far we are only able to take derivatives of explicitly defined functions.

First we have to notice that the point $(\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2})$ is a point of the graph of the function $f_1(x) = \sqrt{1-x^2}$.


We have:

$$f_1'(x) = (\sqrt{1-x^2})' = \frac{1}{2\sqrt{1-x^2}} \cdot (-2x) = \frac{-2x}{2\sqrt{1-x^2}}$$

← chain rule

Hence the slope of the tangent line to the graph of f_1 at the point $x = \frac{\sqrt{2}}{2}$ is:

$$f_1'\left(\frac{\sqrt{2}}{2}\right) = \frac{-2 \cdot \frac{\sqrt{2}}{2}}{2\sqrt{1-\left(\frac{\sqrt{2}}{2}\right)^2}} = \frac{-\sqrt{2}}{2\sqrt{1-\frac{2}{4}}} = \frac{-\sqrt{2}}{2\frac{\sqrt{2}}{2}} = -1$$




and the tangent line has equation:

$$y - \frac{\sqrt{2}}{2} = -1 \left(x - \frac{\sqrt{2}}{2} \right)$$

$$y = -x + \frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}$$

$$y = -x + \sqrt{2}$$



Let us now consider a similar problem with a different curve.

Problem: Find the tangent line to the curve described by the equation:

$$y^3 + xy = \cos(y) - \sin(x)$$

at the point $\left(\frac{\pi}{2}, 0\right)$.

In this case it is not trivial to write y explicitly as a function of x .

Hence for solving the problem we need a new method called **IMPLICIT DIFFERENTIATION**, that allows us to differentiate y with respect to x , without finding an explicit expression for y .

We will describe the method of implicit differentiation in 4 steps:

(1) Take the derivative of each side of the equation with respect to x (do not forget to treat y as a function of x) and apply the differentiation rules

$$\begin{aligned} \text{CHAIN RULE (Leibniz)} \quad \frac{d}{dx} [y^3 + xy] &= \frac{d}{dx} [\cos(y) - \sin(x)] \\ \frac{d}{dx} y^3 &= \frac{d}{dy} y^3 \cdot \frac{dy}{dx} \\ &= 3y^2 \cdot \frac{dy}{dx} \quad \leftarrow \text{chain rule} \\ \frac{d}{dx} (y^3) + \frac{d}{dx} (xy) &= \frac{d}{dx} (\cos(y)) - \frac{d}{dx} (\sin(x)) \\ &= 3y^2 \cdot \frac{dy}{dx} + \left(\frac{d}{dx} (x) \right) \cdot y + x \cdot \frac{dy}{dx} = -\sin(y) \cdot \frac{dy}{dx} - \cos(x) \quad \leftarrow \text{product rule, chain rule} \end{aligned}$$



$$3y^2 \cdot \frac{dy}{dx} + 1 \cdot y + x \cdot \frac{dy}{dx} = -\sin(y) \cdot \frac{dy}{dx} - \cos(x)$$

② You have now an ordinary linear equation where the unknown you want to solve for is $\frac{dy}{dx}$. Solve it

Bring all the terms where $\frac{dy}{dx}$ appears on the same side:

$$\underbrace{3y^2 \frac{dy}{dx} + x \frac{dy}{dx} + \sin(y) \frac{dy}{dx}}_{\text{terms with } \frac{dy}{dx}} = \underbrace{-y - \cos(x)}_{\text{terms without } \frac{dy}{dx}}$$

Factor by $\frac{dy}{dx}$:

$$(3y^2 + x + \sin(y)) \frac{dy}{dx} = -y - \cos(x)$$

Divide by the coefficient of $\frac{dy}{dx}$:

$$\frac{dy}{dx} = -\frac{y + \cos(x)}{3y^2 + x + \sin(y)}$$

note that this time the slope of the tangent line depends on both coordinates of the point.

Note that $\frac{dy}{dx}$ represents the **slope** of the tangent line to the original curve at the generic point (x, y) .

③ Substitute the coordinates of the point in order to find the slope of the tangent line at that point.

$$P\left(\frac{\pi}{2}, 0\right) \Rightarrow \left. \frac{dy}{dx} \right|_{\substack{x=\frac{\pi}{2} \\ y=0}} = -\frac{0 + \cos\left(\frac{\pi}{2}\right)}{3 \cdot 0 + \frac{\pi}{2} + \sin(0)} = \frac{0}{\frac{\pi}{2}} = 0$$

Hence at $P\left(\frac{\pi}{2}, 0\right)$ the tangent line to the curve has slope 0

④ Write an equation of the tangent line

An equation for the tangent line at $P\left(\frac{\pi}{2}, 0\right)$ is $y=0$.

If we go back to the first problem with the circle $x^2 + y^2 = 1$ we see that we get the same result if we apply implicit differentiation:

$$x^2 + y^2 = 1$$

① differentiate both side

$$\frac{d}{dx}(x^2 + y^2) = \frac{d}{dx}(1)$$

$$\frac{d}{dx}(x^2) + \frac{d}{dx}(y^2) = 0$$

② solve for $\frac{dy}{dx}$

$$2x + 2y \cdot \frac{dy}{dx} = 0$$

$$2y \frac{dy}{dx} = -2x$$


$$\frac{dy}{dx} = -\frac{x}{y}$$

③ plug in the coordinates of the point

$$\frac{dy}{dx} \Big|_{\substack{x = \frac{\sqrt{2}}{2} \\ y = \frac{\sqrt{2}}{2}}} = -\frac{\frac{\sqrt{2}}{2}}{\frac{\sqrt{2}}{2}} = -1$$

④ write the equation of the tangent line

$$y - \frac{\sqrt{2}}{2} = -1 \left(x - \frac{\sqrt{2}}{2} \right)$$




Today is **Valentine's day**. Let us make mathematics romantic.

In particular the following equation is very romantic:

$$(x^2 + y^2 - 1)^3 = x^2 y^3$$

Indeed it is the equation of the "**heart curve**":



We want to find the tangent line to the heart at the point $(1, 1)$.

① DIFFERENTIATE BOTH SIDES

$$\frac{d}{dx} (x^2 + y^2 - 1)^3 = \frac{d}{dx} x^2 y^3$$

$$3(x^2 + y^2 - 1)^2 \frac{d}{dx} (x^2 + y^2 - 1) = \left(\frac{d}{dx} (x^2) \right) \cdot y^3 + x^2 \cdot \frac{d}{dx} y^3$$

$$3(x^2 + y^2 - 1)^2 \left[\frac{d}{dx} (x^2) + \frac{d}{dx} (y^2) - \frac{d}{dx} (1) \right] = 2xy^3 + x^2 \cdot 3y^2 \frac{dy}{dx}$$

② SOLVE THE EQUATION FOR $\frac{dy}{dx}$

$$3(x^2 + y^2 - 1)^2 \left[2x + 2y \frac{dy}{dx} \right] = 2xy^3 + x^2 3y^2 \frac{dy}{dx}$$

$$3(x^2 + y^2 - 1)^2 \cdot 2x + 3(x^2 + y^2 - 1)^2 \cdot 2y \frac{dy}{dx} = 2xy^3 + x^2 3y^2 \frac{dy}{dx}$$

$$\left[3(x^2 + y^2 - 1)^2 2y - x^2 3y^2 \right] \frac{dy}{dx} = -3(x^2 + y^2 - 1)^2 2x + 2xy^3$$

$$\frac{dy}{dx} = \frac{-3(x^2 + y^2 - 1)^2 2x + 2xy^3}{3(x^2 + y^2 - 1)^2 2y - x^2 3y^2}$$

③ PLUG IN THE COORDINATES OF THE POINT

$$\left. \frac{dy}{dx} \right|_{x=1, y=1} = \frac{-3(1+1-1)^2 \cdot 2 + 2}{3(1+1-1)^2 \cdot 2 - 3} = \frac{-6 + 2}{6 - 3} = -\frac{4}{3}$$

slope
↓

④ WRITE THE EQUATION OF THE TANGENT LINE

$$y - 1 = -\frac{4}{3}(x - 1) \implies y = -\frac{4}{3}x + \frac{7}{3}$$

