


# INVERSE FUNCTIONS AND LOGARITHMS (Sec. 3.2)

In order to introduce the logarithmic function we need to recall the notion of the inverse of a function when it exists.

Let us consider the following two functions with domain  $\{1, 2, 3, 4\}$  and codomain  $\{a, b, c, d\}$ .


$$f: \{1, 2, 3, 4\} \rightarrow \{a, b, c, d\}$$

$$\begin{aligned} f(1) &= d \\ f(2) &= b \\ f(3) &= a \\ f(4) &= c \end{aligned}$$




$$g: \{1, 2, 3, 4\} \rightarrow \{a, b, c, d\}$$

$$\begin{aligned} g(1) &= b \\ g(2) &= c \\ g(3) &= d \\ g(4) &= a \end{aligned}$$




For each of the previous functions  $f$  and  $g$ , let us see what happens when we exchange the domain with the codomain and we "reverse" the arrows:



This situation corresponds to a new function that we will denote  $f^{-1}$  such that:

$$f^{-1}: \{a, b, c, d\} \rightarrow \{1, 2, 3, 4\}$$

$$\begin{aligned} f^{-1}(a) &= 3 \\ f^{-1}(b) &= 2 \\ f^{-1}(c) &= 4 \\ f^{-1}(d) &= 1 \end{aligned}$$



This situation does not correspond to a function since the input  $b$  gives rise to two outputs 1 and 4.

We say that  $f$  possesses an inverse,  $f^{-1}$ , while  $g$  does not.

If we analyze the situation we remark that for  $f$  all the inputs have a different output, while for  $g$  there are two different values (1 and 4) with the same output ( $b$ ):

$$g(1) = g(4) = b.$$

If  $g^{-1}(b)$  denotes the set of elements of the domain  $\{1, 2, 3, 4\}$  which are sent on  $b$ , we write this fact in the following way:

$$g^{-1}(b) = \{1, 4\}$$

We say that  $f$  is a "one-to-one function" while  $g$  is not.

Def: A function  $f$  is called **one-to-one** if it never takes on the same value twice. In formula:

**[ $p \Rightarrow q$ ]** if  $x_1 \neq x_2 \Rightarrow f(x_1) \neq f(x_2)$  different inputs correspond to different outputs

which is equivalent to:

**[not  $q \Rightarrow$  not  $p$ ]** if  $f(x_1) = f(x_2) \Rightarrow x_1 = x_2$

Remark: If a function  $f: A \rightarrow B$  is one-to-one, then for every element  $b$  in  $B$  there exists at most one element in  $A$  which is sent on  $b$ , i.e.

$f^{-1}(b)$  has at most one element

So, if we go back to the function  $g$ , according to the previous definition, it is not one-to-one because  $g(1) = g(4)$  but  $1 \neq 4$ .


If a function  $f$  is defined on  $\mathbb{R}$  with values in  $\mathbb{R}$ , i.e.  $f: \mathbb{R} \rightarrow \mathbb{R}$ , we can use its graph in the plane for determining whether it is one-to-one.

We have indeed the following geometric method:

### HORIZONTAL LINE TEST

A function is one-to-one if no horizontal line intersects its graph more than once, or, in other words every horizontal line intersects the graph at most once.


ex:  $f(x) = x^2$



X no one-to-one

$f(x_1) = f(x_2)$  with  $x_1 \neq x_2$


$f(x) = e^x$



✓ one to one

every horizontal line intersects the graph AT MOST once

$f(x) = \sin(x)$



X no one-to-one

$f(x_1) = f(x_2) = f(x_3) = \dots$   
with  $x_1, x_2, x_3$  all different each other

Remark: Note that all strictly increasing or strictly decreasing functions are one-to-one.

For one-to-one functions we can define an inverse function:

Def: Let  $f$  be a one-to-one function with domain  $A$  and range  $B$ .


Then its **inverse function**  $f^{-1}$  has domain  $B$  and range  $A$  and it is defined by:

for all  $y$  in  $B$ ,  $f^{-1}(y) = x \iff f(x) = y$ .

If  $f: A \rightarrow B$  and  $f^{-1}: B \rightarrow A$  is its inverse function then  $f$  and  $f^{-1}$  verify the following **cancellation equations**:


for all  $x$  in  $A$

$f^{-1}(f(x)) = x$



for all  $y$  in  $B$

$f(f^{-1}(y)) = y$



Remark: Note that  $f^{-1}(x) \neq [f(x)]^{-1} !!$

inverse function of  $f$

reciprocal function of  $f: \frac{1}{f(x)}$

ex:  $f = e^x$  is a one to one function with domain  $\mathbb{R}$  and range  $(0, \infty)$ :

$$f: \mathbb{R} \rightarrow (0, \infty)$$

$$\begin{array}{l} 0 \mapsto 1 \\ 1 \mapsto e \\ \vdots \\ \text{etc.} \end{array}$$

Hence  $f$  possesses an inverse  $f^{-1}$  with domain  $(0, \infty)$  and range  $\mathbb{R}$ :


$$f^{-1}: (0, \infty) \rightarrow \mathbb{R}$$

$$\begin{array}{l} 1 \mapsto 0 \\ e \mapsto 1 \end{array}$$

$$\begin{array}{l} f^{-1}(1) = 0 \text{ since } f(0) = 1 \\ f^{-1}(e) = 1 \text{ since } f(1) = e \end{array}$$

EXERCISE: Find the inverse function of  $f(x) = x^3 + 1$ .


First of all  $f$  is one-to-one, since it passes the horizontal line test.



If  $f(x) = y$  then we define  $f^{-1}(y) = x$ .

$$y = x^3 + 1 \Rightarrow x^3 = y - 1 \Rightarrow x = \sqrt[3]{y - 1}$$

$$\text{Then } f^{-1}(y) = x = \sqrt[3]{y - 1}$$



Remark that this graph is obtained by reflecting the graph of  $f$  about the line  $y = x$ .

Theorem: If  $f$  is a one-to-one continuous function defined on an interval, then its inverse  $f^{-1}$  is also continuous

## LOGARITHMIC FUNCTION

For  $a > 0$ ,  $a \neq 1$  the exponential function  $f(x) = a^x$  is strictly increasing or decreasing. It is then one-to-one.

Its inverse  $f^{-1}$  is called logarithmic function with base  $a$  and denoted  $\log_a$ .

By definition we have

$$\log_a x = y \iff a^y = x$$


From the properties of the exponential function we can deduce properties for the logarithmic function

### PROPERTIES OF $a^x$

- domain  $\mathbb{R}$
- range  $(0, \infty)$
- continuous

### PROPERTIES OF $\log_a x$


- range  $\mathbb{R}$
- domain  $(0, \infty)$
- continuous



$$0 < a < 1$$

$$\lim_{x \rightarrow 0^+} \log_a x = \infty$$

$$\lim_{x \rightarrow \infty} \log_a x = -\infty$$



$$a > 1$$

$$\lim_{x \rightarrow 0^+} \log_a x = -\infty$$

$$\lim_{x \rightarrow \infty} \log_a x = \infty$$

Cancellation laws:  $f(x) = a^x$ ,  $f^{-1}(x) = \log_a(x)$

For all  $x$  in  $\mathbb{R}$ ,  $f^{-1}(f(x)) = x \rightsquigarrow \log_a(a^x) = x$ , for all  $x$  in  $\mathbb{R}$

For all  $x$  in  $(0, \infty)$ ,  $f(f^{-1}(x)) = x \rightsquigarrow a^{\log_a x} = x$ , for all  $x > 0$ .

ex:  $\log_2(8) = 3$  since  $2^3 = 8$ .

•  $\log_a 1 = 0$  since  $a^0 = 1$ , for all  $a > 0$ ,  $a \neq 1$ .

The laws of exponential can be turned in laws of logarithms:

### LAWS OF LOGARITHMS

If  $a > 0$ ,  $a \neq 1$  and  $x, y > 0$  then

$$(1) \log_a(xy) = \log_a(x) + \log_a(y)$$

$$(2) \log_a\left(\frac{x}{y}\right) = \log_a(x) - \log_a(y)$$

$$(3) \log_a(x^r) = r \log_a(x), \text{ where } r \text{ is any real number}$$

Proof

$$(1) a^{\log_a(x) + \log_a(y)} = a^{\log_a(x)} \cdot a^{\log_a(y)} = x \cdot y \Rightarrow$$

$a^{x+y} = a^x \cdot a^y$        $a^{\log_a(x)} = x$

$$\Rightarrow \text{by definition } \log_a(xy) = \log_a(x) + \log_a(y)$$

$$(2) a^{\log_a(x) - \log_a(y)} = \frac{a^{\log_a(x)}}{a^{\log_a(y)}} = \frac{x}{y}$$

$a^{x-y} = \frac{a^x}{a^y}$        $a^{\log_a(x)} = x$

$$\Rightarrow \text{by definition } \log_a\left(\frac{x}{y}\right) = \log_a(x) - \log_a(y)$$

$$(3) a^{r \log_a(x)} = \left(a^{\log_a(x^r)}\right) = x^r$$

$a^{xy} = (a^x)^y$        $a^{\log_a(x^r)} = x^r$

$$\Rightarrow \text{by definition } \log_a(x^r) = r \log_a(x)$$

ex:  $\log_3 18 - \log_3 2 = \log_3 \frac{18}{2} = \log_3 9 = \log_3 3^2 = 2.$

As in the case of the exponential function, there exists a "most convenient" base for the logarithm function, which is again the number  $e$ .

### Notation

$$\log x = \log_{10} x$$

$$\ln x = \log_e x$$

## Natural logarithm

The logarithm function with base  $e$  is called **natural logarithm** and is denoted:

$$f(x) = \ln(x)$$

### Properties of $\ln(x)$

- domain  $(0, \infty)$
- range  $\mathbb{R}$
- continuous on  $(0, \infty)$
- $\lim_{x \rightarrow 0^+} \ln(x) = -\infty$
- $\lim_{x \rightarrow \infty} \ln(x) = \infty$
- for all  $x > 0$   $\ln x = y \Leftrightarrow e^y = x$
- $\ln(e^x) = x$  for all  $x$  in  $\mathbb{R}$
- $e^{\ln(x)} = x$  for all  $x$  in  $\mathbb{R}$
- $\ln 1 = 0$
- $\ln e = 1$
- $\log_a x = \frac{\ln(x)}{\ln(a)}$

